

Reconstructing a Simple Polytope from its Graph

Volker Kaibel*

TU Berlin, MA 6-2
Straße des 17. Juni 136
10623 Berlin, Germany

kaibel@math.tu-berlin.de

<http://www.math.tu-berlin.de/~kaibel>

Abstract. Blind and Mani [2] proved that the entire combinatorial structure (the vertex-facet incidences) of a simple convex polytope is determined by its abstract graph. Their proof is not constructive. Kalai [15] found a short, elegant, and algorithmic proof of that result. However, his algorithm has always exponential running time. We show that the problem to reconstruct the vertex-facet incidences of a simple polytope P from its graph can be formulated as a combinatorial optimization problem that is strongly dual to the problem of finding an abstract objective function on P (i.e., a shelling order of the facets of the dual polytope of P). Thereby, we derive polynomial certificates for both the vertex-facet incidences as well as for the abstract objective functions in terms of the graph of P . The paper is a variation on joint work with Michael Joswig and Friederike Körner [12].

1 Introduction

The *face lattice* L_P of a (convex) polytope P is any lattice that is isomorphic to the lattice formed by the set of all faces of P (including $\emptyset$ and P itself), ordered by inclusion. It is well-known to be determined by the *vertex-facet incidences* of P , i.e., by any graph that is isomorphic to the bipartite graph whose nodes are the vertices and the facets of P , where the edges are defined by the pairs $\{v, f\}$ of vertices v and facets f with $v \in f$. In lattice theoretic terms, L_P is a ranked, atomic, and coatonic lattice, and thus, the sub-poset formed by its atoms and coatoms already determines the whole lattice. Actually, one can compute L_P from the vertex-facet incidences of P in $\mathcal{O}(\eta \cdot \alpha \cdot \lambda)$ time, where η is the minimum of the number of vertices and the number of facets, α is the number of vertex-facet incidences, and λ is the total number of faces of P [14].

The *graph* $G_P = (V_P, E_P)$ of a polytope P is any graph that is isomorphic to the graph whose nodes are the vertices of P , where two nodes are adjacent if and only if the convex hull of the corresponding two vertices is a one-dimensional face of P . Phrased differently, G_P is the graph defined on the rank one elements of L_P ,

* Supported by the Deutsche Forschungsgemeinschaft, FOR 413/1-1 (Zi 475/3-1).

where two rank one elements are adjacent if and only if they are below a common rank two element. While the vertex-facet incidences completely determine the face-lattice of any polytope, the graph of a polytope in general does not encode the entire combinatorial structure. This can be seen, e.g., from the examples of the cut polytope associated with the complete graph on n nodes and the $\binom{n}{2}$ -dimensional cyclic polytope with 2^{n-1} vertices, which both have complete graphs. Another example is the four-dimensional polytope shown in Fig. 1 whose graph is isomorphic to the graph of the five-dimensional cube.




Fig. 1. A Schlegel-diagram (projection onto one facet) of a four-dimensional polytope with the graph of a five-dimensional cube, found by Joswig & Ziegler [13].

Actually, in all dimensions less than four such ambiguities cannot occur. For one- or two dimensional polytopes this is obvious, and for three-dimensional polytopes it follows from Whitney’s theorem [18] saying that every 3-connected planar graph has a unique (up to reflection) plane embedding.

A d -dimensional polytope P is *simple* if every vertex of P is contained in precisely d facets, which is equivalent to G_P being d -regular (the polytope is non-degenerate in terms of Linear Programming). Every face of a simple polytope is simple as well. None of the examples showing that it is, in general, impossible to reconstruct the face lattice of a polytope from its graph, is simple.

In fact, Blind and Mani [2] proved in 1987 that the face lattice of a simple polytope is determined by its graph. Their proof (which we sketch in Sect. 2) is not constructive and crucially relies on the topological concept of homology. In 1988, Kalai [15] found a short and elegant proof (reviewed in Sect. 3) that does only use elementary geometric and combinatorial reasoning with the main advantage of being algorithmic. However, the running time of the method that can be devised from it is exponential in the size of the graph.

Perles conjectured in the 1970’s (see [15]) that for a d -dimensional simple polytope P every subset $F \subset V_P$ that induces a $(d - 1)$ -regular, connected, and non-separating subgraph of G_P corresponds to the vertex set of a facet of P . A proof of this conjecture would have lead immediately to a polynomial time

algorithm that, given the graph $G_P = (V_P, E_P)$ of a simple polytope P , decides for a set of subsets of V_P if it corresponds to the set of vertex sets of facets of P . However, Haase and Ziegler [10] recently disproved Perles' conjecture. They found a four-dimensional simple polytope whose graph has a 3-regular, non-separating, and even 3-connected induced subgraph that does not correspond to any facet.

Refining ideas from Kalai's proof (Sect. 4), we show that the problem of reconstructing the vertex-facet incidences of a simple polytope P from its graph G_P can be formulated as a combinatorial optimization problem that has a well-stated strongly dual problem (Sect. 5). The optimal solutions to this dual problem are certain orientations of G_P (induced by "abstract objective functions") that are important also in different contexts. In particular, we provide short certificates for both the vertex-facet incidences of a simple polytope and for the abstract objective functions in terms of G_P . We conclude with some remarks on the complexity status of the problem to decide whether a claimed solution to the reconstruction problem indeed are the vertex-facet incidences of the respective polytope in Sect. 6.

The material presented here has evolved from joint work with Michael Joswig and Friederike Körner [12]. The basic ideas and results are the same in both papers. However, the concept of a "facoidal system of walks" is newly introduced here. It differs from the corresponding notion of a "2-system" (introduced in [12]) with the main effect that one knows how to compute efficiently some facoidal system of walks from the graph G_P of a simple polytope P (see Proposition 3), while it is unclear how to find a 2-system from G_P efficiently. Furthermore, the proof of Theorem 3 we give here is different from the corresponding proof in [12]. Finally, the complexity theoretic statement of Corollary 5 does not appear in [12].

For all notions from the theory of polytopes that we use without (sufficient) explanations, we refer to Ziegler's book [19]. We use the terms *d-polytope* and *k-face* for d -dimensional polytopes and k -dimensional faces, respectively. Often we will identify a face F of a simple polytope P with the subset of nodes of G_P that corresponds to the vertex set of F . Whenever we talk about "polynomial time" or "efficient" this refers to the size of the graph G_P of the respective (simple) polytope P .

2 The Theorem of Blind and Mani

Blind and Mani [2] proved their theorem in the dual setting, i.e., for simplicial rather than for simple polytopes. Nevertheless, we sketch parts of their proof in terms of simple polytopes here. The starting point is the observation that, while a priori it is by no means clear if the graph $G_P = (V_P, E_P)$ of a simple polytope P determines the face lattice of P , it is easy to see that the 2-faces of P (as subsets of V_P) carry the entire information on the combinatorial structure of P .

Let P be a simple polytope. For a node $v \in V_P$ of G_P denote by $\Gamma(v) \subset V_P$ the subset of nodes that are adjacent to v (the *neighbors* of v). For any k -element

subset $S \subset \Gamma(v)$ there is a k -face of P that contains the vertices that correspond to $S \cup \{v\}$ (and no vertex that corresponds to a node in $\Gamma(v) \setminus S$). We call the subset $F(S \cup \{v\}) \subseteq V_P$ of nodes corresponding to the vertices of that face the k -face spanned by $S \cup \{v\}$.

For an edge $e = \{v, w\} \in E_P$ let $\Psi_{(v,w)} : \Gamma(v) \setminus \{w\} \longrightarrow \Gamma(w) \setminus \{v\}$ be the map that assigns to each subset $S \subset \Gamma(v) \setminus \{w\}$ the subset $T \subset \Gamma(w) \setminus \{v\}$ with $F(S \cup \{v, w\}) = F(T \cup \{w, v\})$. The maps $\Psi_{(v,w)}$ are cardinality preserving bijections, where $\Psi_{(w,v)}$ is the inverse of $\Psi_{(v,w)}$.

Proposition 1. *Let P be a simple polytope. For each $e = \{v, w\} \in E_P$ and $S \subseteq \Gamma(v) \setminus \{w\}$ we have*

$$\Psi_{(v,w)}(S) = \overline{\Psi_{(v,w)}(\overline{S})}$$

(where $\overline{U}$ is the respective complement of the set U).

Proof. This follows from the fact that we have $\Psi_{(v,w)}(S_1 \cap S_2) = \Psi_{(v,w)}(S_1) \cap \Psi_{(v,w)}(S_2)$ for all $S_1, S_2 \subseteq \Gamma(v) \setminus \{w\}$.

With the notations of Proposition 1 denote by $\Psi_{(v,w)}^k$ the restriction of the map $\Psi_{(v,w)}$ to the $(k-1)$ -element subsets of $\Gamma(v) \setminus \{w\}$. There are quite obvious algorithms that compute from the maps $\Psi_{(v,w)}^k, \{v, w\} \in E_P$, the k -faces of P (as subsets of V_P), and vice versa, in polynomial time in the number $f_k(P)$ of k -faces of the simple d -polytope P . Since both $f_2(P)$ as well as $f_{d-1}(P)$ are bounded polynomially in the size of G_P , the following result follows.

Corollary 1. *There are polynomial time algorithms that, given the graph G_P of a simple d -polytope P , compute the set of facets of P from the set of 2-faces of P (both viewed as sets of subsets of V_P), and vice versa.*

For the rest of this section let P_1 and P_2 be two simple polytopes, and let $g : V_{P_1} \longrightarrow V_{P_2}$ be an isomorphism of the graphs $G_{P_1} = (V_{P_1}, E_{P_1})$ and $G_{P_2} = (V_{P_2}, E_{P_2})$ of P_1 and P_2 , respectively (i.e., g is an in both directions edge preserving bijection).

The core of Blind and Mani's paper [2] is the following result.

Proposition 2. *The graph isomorphism g maps every cycle in G_{P_1} that corresponds to a 2-face of P_1 to a cycle in G_{P_2} that corresponds to a 2-face of P_2 .*

Blind and Mani's proof proceeds in the dual setting, i.e., in terms of the boundary complexes ∂P_1^* and ∂P_2^* of the simplicial dual polytopes P_1^* and P_2^* of P_1 and P_2 , respectively. The strategy is to show that, if some cycle in G_{P_1} corresponding to a 2-face of P_1 was mapped to some cycle in G_{P_2} that does not correspond to any 2-face of P_2 , then a certain sub-complex of ∂P_2^* would have a certain non-vanishing (reduced) homology group. They complete their proof of Proposition 2 by showing that the respective homology group, however, is zero. The key ingredient they use to prove this is the following. For each face F of P_1^*

there is a *shelling order* of the facets of P_1^* (i.e., an ordering satisfying certain convenient topological properties, which, however, can be expressed completely combinatorially, see Sect. 3) in which the facets containing F come first.

From Proposition 2 one can deduce that the graph isomorphism g actually induces a bijection between the cycles in G_{P_1} that correspond to 2-faces of P_1 and the cycles in G_{P_2} that correspond to 2-faces of P_2 . Once this is established, Proposition 1 yields the following result.

Theorem 1 (Blind & Mani [2]). *Every isomorphism between the graphs G_{P_1} and G_{P_2} of two simple polytopes P_1 and P_2 , respectively, induces an isomorphism between the vertex-facet incidences of P_1 and P_2 . In particular, the graph of a simple polytope determines its entire face lattice.*

3 Kalai's Constructive Proof

Kalai realized that the existence of shelling orders as exploited by Blind and Mani can be used directly in order to devise a simple proof which does not rely on any topological notions like homology [15]. He formulated his proof in the original setting, i.e., for simple polytopes, where the notion corresponding to “shelling” is called “abstract objective function.”

From now on, let P be a simple d -polytope with n vertices. For simplicity of notation, we will identify each face of P not only with the corresponding subset of V_P , but also with the corresponding induced subgraph of G_P . Furthermore, by saying that $w \in W \subset V_P$ is a sink of W we mean that w is a sink of the orientation induced on the subgraph of G_P that is induced by W .

Definition 1. *Every bijection $\varphi : V_P \longrightarrow \{1, \dots, n\}$ induces an acyclic orientation $\mathcal{O}_\varphi$ of the graph G_P of P , where an edge is directed from its larger end-node to its smaller end-node (with respect to φ). The map φ is called an abstract objective function (AOF) if $\mathcal{O}_\varphi$ has a unique sink in every non-empty face of P (including P itself). Such an orientation of G_P is called an AOF-orientation.*

The inverse orientation of an AOF-orientation is an AOF-orientation as well (this follows, e.g., from Theorem 3). Thus, every AOF-orientation also has a unique source in every non-empty face.

From the fact that the simplex algorithm works correctly (on every face) one easily derives that every linear function that assigns pairwise different values to the vertices of P induces an AOF-orientation (this is a consequence of the convexity of the faces). From this observation, the following fact follows (which is dual to the existence of the shelling orders required in Blind and Mani's proof).

Lemma 1. *Let $W \subset V_P$ be any face of P . There is an AOF-orientation of G_P for which W is terminal, i.e., no edge in the cut defined by W is directed from W to $V_P \setminus W$.*

In a sense, this statement can be reversed.

Lemma 2. *Let $W \subset V_P$ be a set of nodes inducing a k -regular connected subgraph of G_P , and let $\mathcal{O}$ be an AOF-orientation for which W is terminal. Then W is a k -face of P .*

Proof. Since $\mathcal{O}$ is acyclic, it has a source s in W . Let $w_1, \dots, w_k \in W$ be the neighbors of s in W , and let $F := F(\{s, w_1, \dots, w_k\}) \subset V_P$ be the k -face of P that is spanned by $s, w_1, \dots, w_k$. Since $\mathcal{O}$ has unique sources on non-empty faces, $s \in W \cap F$ must be the unique source of F . By the acyclicity of $\mathcal{O}$ there hence is a monotone path from s to every node in F . Since W is terminal this implies $F \subseteq W$. Because both F and W induce k -regular connected subgraphs of G_P , $F = W$ follows.

Lemma 1 and Lemma 2 imply that one can compute the vertex-facet incidences of P , provided that one knows all AOF-orientations of G_P . Kalai's crucial discovery is that one can compute the AOF-orientations just from G_P (i.e., without explicitly knowing the faces of P).

Definition 2. *For an orientation $\mathcal{O}$ of G_P let $h_k(\mathcal{O})$ be the number of nodes with in-degree k . The number*

$$\mathcal{H}(\mathcal{O}) := \sum_{k=0}^d h_k(\mathcal{O}) \cdot 2^k$$

is called the $\mathcal{H}$ -sum of $\mathcal{O}$.

Since every subset of neighbors of a vertex v of P together with v span a face of P containing no other neighbors of v , one finds (by double-counting) that

$$\mathcal{H}(\mathcal{O}) = \sum_{F \text{ face of } P} (\# \text{ sinks of } \mathcal{O} \text{ in } F) \quad (1)$$

is the total number of sinks induced by $\mathcal{O}$ on faces of P . Consequently, since every acyclic orientation has at least one sink in every non-empty face, we have the following characterization.

Lemma 3. *An orientation $\mathcal{O}$ of G_P is an AOF-orientation if and only if it is acyclic and has minimal $\mathcal{H}$ -sum among all acyclic orientations of G_P (which then equals the number of non-empty faces of P).*

Thus, by enumerating all $2^{\frac{d+n}{2}} = \sqrt{2}^{d \cdot n}$ orientations of G_P one can find all AOF-orientations of G_P .

Theorem 2 (Kalai [15]). *There is an algorithm that computes the vertex-facet incidences of a simple d -polytope with n vertices from its graph in $\mathcal{O}(\sqrt{2}^{d \cdot n})$ steps.*

4 Walks and Orientations

In this section, we refine the ideas of Kalai's proof and combine them with the observation (exploited by Blind and Mani) that it suffices to identify the 2-faces from the graph of a simple polytope, even with respect to the question for polynomial time reconstruction algorithms (see Corollary 1). Let us start with a result that emphasizes the importance of the 2-faces even more. The result was known for cubes [11]; for three-dimensional simple polytopes it was independently proved by Develin [4]). For general simple polytopes it seems that it was assumed to be false (see [19, Ex. 8.12 (iv)]).

Theorem 3. *An acyclic orientation $\mathcal{O}$ of the graph G_P of a simple polytope P is an AOF-orientation if and only if it has a unique sink on every 2-face of P .*

Proof. The “only if” part is clear by definition. For the “if” part, let $\varphi : V_P \rightarrow \{1, \dots, n\}$ be a bijection inducing an acyclic orientation $\mathcal{O} = \mathcal{O}_\varphi$ that has a unique sink in every 2-face of P . Suppose there is a face F of P in which $\mathcal{O}$ has two sinks $t_1, t_2 \in F$ ($t_1 \neq t_2$). We might assume $F = P$ (because F itself is a simple polytope with every 2-face of F being a 2-face of P).

Since G_P is connected, there is a path in G_P connecting t_1 and t_2 . Let $\Pi \neq \emptyset$ be the set of all these paths. For every $\pi \in \Pi$ we denote by $\mu(\pi)$ the maximal φ -value of any node in π . Let $\pi_{\min} \in \Pi$ be a path with minimal μ -value among all paths in Π , where $v \in V_P$ is the node in $\pi_{\min}$ with $\varphi(v) = \mu(\pi_{\min})$ (see Fig. 2).




Fig. 2. Illustration of the proof of Theorem 3. The fat grey path is the one yielding the contradiction.

Obviously, v is a source in the path $\pi_{\min}$ (in particular, $v \notin \{t_1, t_2\}$). Let C be the 2-face spanned by v and its two neighbors v_1 and v_2 in $\pi_{\min}$. Since v is the unique source of $\mathcal{O}$ in C , v has the largest φ -value among all nodes in the union U of C and $\pi_{\min}$. But $U \setminus \{v\}$ induces a connected subgraph of G_P containing both t_1 and t_2 , which contradicts the minimality of $\mu(\pi_{\min})$.

From now on let, again, P be a simple polytope. The ultimate goal is to find the system of cycles in the graph G_P that corresponds to the set of 2-faces

of P . However, we even do not know how to *prove* or *disprove* efficiently that a given system of cycles actually is the one we are searching for. We now define more general systems having the property that one can at least generate one of them in polynomial time (which in general, of course, will not be the desired one), and among which the one corresponding to the set of 2-faces of P can be characterized using AOF-orientations.

- Definition 3.** 1. A sequence $W = (w_0, \dots, w_{l-1})$ (with $l \geq 3$) of nodes in G_P is called a *closed smooth walk* in G_P , if $\{w_i, w_{i+1}\}$ is an edge of G_P and $w_{i-1} \neq w_{i+1}$ for all i (where, as in the following, all indices are taken modulo l). Note that the w_i need not be pairwise disjoint. We will identify two closed smooth walks if they differ only by a cyclic shift and/or a “reflection” of their node sequences.
2. A set $\mathcal{W}$ of closed smooth walks in G_P is a *facoidal system* of walks if for every triple $v, v_1, v_2 \in V_P$ ($v_1 \neq v_2$) such that both v_1 and v_2 are neighbors of v there is a unique closed smooth walk $(w_0, \dots, w_{l-1}) \in \mathcal{W}$ with $(w_{i-1}, w_i, w_{i+1}) \in \{(v_1, v, v_2), (v_2, v, v_1)\}$ for some i , which is also required to be unique.

The system of 2-faces of P yields a uniquely determined (recall the identifications mentioned in part 1 of the definition) facoidal system of walks in G_P , which is denoted by $\mathcal{C}_P$. In general, there are many other facoidal systems of walks (see Fig. 3).




Fig. 3. A facoidal system of four walks in the graph of the three-dimensional cube.

For each path λ in G_P of length two denote by $v(\lambda)$ the inner node of λ . Let G_P^* be the graph defined on the paths of length two in G_P , where two paths λ_1 and λ_2 are adjacent if and only if they share a common edge and $v(\lambda_1) \neq v(\lambda_2)$ holds (see Fig. 4).

A *2-factor* in a graph G is a set of (not self-intersecting) cycles in G such that every node is contained in a unique cycle. Checking whether a graph has a 2-factor and finding one (if it exists) can be reduced (by a procedure due to




Fig. 4. The left constellation gives rise to an edge of G_P^* , while the right one does not.

Tutte [17]) to searching for a perfect matching in a related graph (which can be performed in polynomial time by Edmonds' algorithm [6]).

Proposition 3. *For simple polytopes P ,*

1. *there is a (polynomial time computable) bijection between the facoidal systems of walks in G_P and the 2-factors of G_P^* ,*
2. *checking whether a given set of node-sequences in G_P is a facoidal system of walks can be done in polynomial time, and*
3. *one can find a facoidal system of walks in G_P in polynomial time.*

Proof. Part 2 is obvious, part 3 follows from part 1 by Tutte's reduction [17] and Edmonds algorithm [6], and part 1 is readily obtained from the definitions.

Proposition 3 shows that facoidal systems of walks have quite convenient algorithmic properties. However, they become useful only due to the fact that the system $\mathcal{C}_P$ corresponding to the 2-faces of P can be well-characterized among them, as we will demonstrate next.

Definition 4. *Let $\mathcal{O}$ be any orientation of G_P .*

1. *The $\mathcal{H}_2$ -sum of $\mathcal{O}$ is defined as*

$$\mathcal{H}_2(\mathcal{O}) := \sum_{k=0}^d h_k(\mathcal{O}) \cdot \binom{k}{2}.$$

2. *A closed smooth walk $(w_0, \dots, w_{l-1})$ in G_P has a sink (source, respectively) at position i (with respect to the orientation $\mathcal{O}$), if the edges $\{w_i, w_{i-1}\}$ and $\{w_i, w_{i+1}\}$ both are directed towards (away from, respectively) w_i .*

The following follows immediately from the definitions.

Lemma 4. *For every orientation $\mathcal{O}$ of G_P the sum $\mathcal{H}_2(\mathcal{O})$ equals the total number of sinks (with respect to $\mathcal{O}$) in every facoidal system of walks in G_P .*

Now we can formulate and prove the main result of this section (where $f_2(P)$ denotes the number of 2-faces of P).

Theorem 4. *Let P be a simple polytope, $\mathcal{W}$ a facoidal system of walks in G_P , and $\mathcal{O}$ an acyclic orientation of G_P . Then*

$$\#\mathcal{W} \leq f_2(P) \leq \mathcal{H}_2(\mathcal{O}) \quad (2)$$

holds.

1. *The first inequality holds with equality if and only if $\mathcal{W} = \mathcal{C}_P$ (i.e., $\mathcal{W}$ “is” the set of 2-faces of P).*
2. *The second inequality holds with equality if and only if $\mathcal{O}$ is an AOF-orientation of G_P .*

Proof. Since $\mathcal{O}$ is acyclic, every closed smooth walk in G_P must have at least one sink with respect to $\mathcal{O}$. Thus, Lemma 4 implies

$$\#\mathcal{W} \leq \mathcal{H}_2(\mathcal{O}) \quad , \quad (3)$$

yielding

$$f_2(P) = \#\mathcal{C}_P \leq \mathcal{H}_2(\mathcal{O}) \quad , \quad (4)$$

where by Theorem 3 equality holds in (4) if and only if $\mathcal{O}$ is an AOF-orientation. Because G_P has an AOF-orientation $\mathcal{O}_0$ (see Lemma 1), inequality (3) gives $\#\mathcal{W} \leq \mathcal{H}_2(\mathcal{O}_0) = f_2(P)$.

Hence, it remains to prove that $\#\mathcal{W} = \#\mathcal{C}_P$ implies $\mathcal{W} = \mathcal{C}_P$. Suppose, that $\#\mathcal{W} = \#\mathcal{C}_P$ holds. It thus suffices to show $\mathcal{C}_P \subseteq \mathcal{W}$ (since we know already $\#\mathcal{C}_P \geq \#\mathcal{W}$). Let $C \in \mathcal{C}_P$ be any closed smooth walk corresponding to a 2-face of P . By Lemma 2 there is an AOF-orientation $\mathcal{O}_C$ of G_P such that C is terminal with respect to $\mathcal{O}_C$. Let $w_1 \in V_P$ be the unique source in C (with respect to $\mathcal{O}_C$), and let w_0 and w_2 be the two neighbors of w_1 in C . By definition, there is a (unique) $W = (w_0, w_1, w_2, \dots, w_{l-1}) \in \mathcal{W}$. Because of $\#\mathcal{W} = \#\mathcal{C}_P = \mathcal{H}_2(\mathcal{O}_C)$ the closed smooth walk W has a unique sink at some position j and its unique source at position 1. Thus, the two paths $(w_1, w_2, \dots, w_j)$ and $(w_1, w_0, \dots, w_j)$ both are monotone. Since C is terminal, this implies that these two paths are contained in C . Therefore we have $C = W \in \mathcal{W}$.

5 Good Characterizations

Theorem 4 immediately yields characterizations of sets of 2-faces and of AOF-orientations that are similar to Kalai’s characterization of AOF-orientations (see Lemma 3).

Corollary 2. *Let P be a simple polytope.*

1. *A facoidal system of walks in G_P is the system $\mathcal{C}_P$ of 2-faces of P if and only if it has maximal cardinality among all facoidal systems of walks in G_P .*
2. *An acyclic orientation of G_P is an AOF-orientation if and only if it has minimal $\mathcal{H}_2$ -sum among all acyclic orientations of G_P .*




Fig. 5. Illustration of the proof of Theorem 4. If $W \neq C$ then C cannot be terminal.

Unfortunately, for arbitrary graphs the problem of finding a 2-factor with as many cycles as possible is $\mathcal{NP}$ -hard. This follows from the fact that the question whether a graph can be partitioned into triangles is $\mathcal{NP}$ -complete [7, Prob. GT11].

With respect to algorithmic questions, the following *good characterizations* (in the sense of Edmonds [5,6]) of the set of 2-faces (and thus, by Proposition 1 of the vertex-facet incidences) as well as of AOF-orientations may be more valuable than those in Corollary 2.

Corollary 3. *Let P be a simple polytope.*

1. *Let $\mathcal{W}$ be a facoidal system of walks in G_P . Either there is an acyclic orientation of G_P having a unique sink in every walk of $\mathcal{W}$, or there is a facoidal system of walks in G_P of larger cardinality than $\#\mathcal{W}$. In the first case, $\mathcal{W} = \mathcal{C}_P$ “is” the set of 2-faces of P , in the second, it is not.*
2. *Let $\mathcal{O}$ be an acyclic orientation of G_P . Either there is a facoidal system $\mathcal{W}$ of walks in G_P such that $\mathcal{O}$ has a unique sink in every walk in $\mathcal{W}$, or there is an acyclic orientation of G_P with smaller $\mathcal{H}_2$ -sum than $\mathcal{H}_2(\mathcal{O})$. In the first case, $\mathcal{O}$ is an AOF-orientation, in the second, it is not.*

For graphs G of simple polytopes let us define *Problem (A)* as

$$\max \#\mathcal{W} \quad \text{subject to} \quad \mathcal{W} \text{ facoidal system of walks in } G$$

and *Problem (B)* as

$$\min \mathcal{H}_2(\mathcal{O}) \quad \text{subject to} \quad \mathcal{O} \text{ acyclic orientation of } G.$$

A third consequence of Theorem 4 is the following result.

Corollary 4. *The Problems (A) and (B) form a pair of strongly dual combinatorial optimization problems. The optimal solution of Problem (A) yields the 2-faces of the respective polytope (and thus its vertex-facet incidences, see Proposition 1). Every optimal solution to Problem (B) is an AOF-orientation of the graph.*

Thus, the answer to Perles' original question whether the vertex-facet incidences of a simple polytope are at all determined by its graph, is not only “yes” (as proved by Blind and Mani), or “yes, and they can be computed” (as shown by Kalai), but at least “yes, and they can be computed by solving a combinatorial optimization problem that has a well-stated strongly dual problem.”

6 Remarks

Corollary 4 suggests to design a primal-dual algorithm for the problem of reconstructing (the vertex-facet incidences of) a simple polytope from its graph. Such an algorithm would start by computing an arbitrary facoidal system $\mathcal{W}$ of walks in the given graph (see Proposition 3) and any acyclic orientation $\mathcal{O}$. Then it would check for $\#\mathcal{W} = \mathcal{H}_2(\mathcal{O})$. If equality holds then by Theorem 4 one is done. Otherwise, the algorithm would try to improve either $\mathcal{W}$ or $\mathcal{O}$ by exploiting the reasons for $\#\mathcal{W} \neq \mathcal{H}_2(\mathcal{O})$. For a concise treatment of different classical and recent applications of the primal-dual method in Combinatorial Optimization see [8].

Such a (polynomial time) primal-dual algorithm would in particular yield polynomial time algorithms for the problem to determine an (arbitrary) AOF-orientation from the graph G_P of a simple polytope P and the set of 2-faces of P , as well as for the problem to determine the 2-faces of P from G_P and an AOF-orientation. As for the first of these two problems it is worth to mention that no polynomial time method is known that would find any AOF-orientation even if the input is the entire face lattice of P . For the second problem no polynomial time algorithm is known as well.

Let (C) be the problem to decide for the graph G_P of a simple polytope P and a set $\mathcal{C}$ of subsets of nodes of G_P if $\mathcal{C}$ is the set of the subsets of nodes of G_P that correspond to the 2-faces of P . Let (D) be the problem to decide for the graph G_P of a simple polytope P and an orientation $\mathcal{O}$ of G_P if $\mathcal{O}$ is an AOF-orientation. The good characterizations in Corollary 3 may tempt one to conjecture that these two problems can be solved in polynomial time.

Unfortunately, from the complexity theoretic point of view, Corollary 3 does not provide us with any evidence for that. In particular, it does *not* imply that problems (C) and (D) are contained in $\mathcal{NP} \cap \text{co}\mathcal{NP}$. The reason is that the problem (G) to decide for a given graph if there is any simple polytope P such that G is isomorphic to the graph of P is neither known to be in $\mathcal{NP}$ nor in $\text{co}\mathcal{NP}$. Corollary 3 only shows that both problems (C) and (D) are in $\mathcal{NP} \cap \text{co}\mathcal{NP}$, if one restricts them to any class of graphs for which problem (G) is in $\mathcal{NP} \cap \text{co}\mathcal{NP}$.

The problem (S) to decide for a given lattice L if there is a simple polytope P such that the face lattice L_P of P is isomorphic to L is known as the *Steinitz problem for simple polytopes*.

Corollary 5. *If problem (G) is contained in $\mathcal{NP}$ or $\text{co}\mathcal{NP}$, then problem (S) is contained in $\mathcal{NP}$ or $\text{co}\mathcal{NP}$, respectively.*

Proof. The face lattice L_P of a polytope P is ranked. The graph G having the rank one elements of L_P as its nodes, where two rank one elements are adjacent if and only if they are below a common rank two element, is isomorphic to the graph of P . It can be computed from L_P in polynomial time. Corollary 1 shows that one can compute the vertex-facet incidences (and thus the entire face lattice L_P , see the first paragraph of the introduction) of a simple polytope P in polynomial time (in the size of L_P) from the poset that is induced by the elements of rank one (corresponding to the vertices), rank two (corresponding to the 1-faces), and rank three (corresponding to the 2-faces). Together with the first part of Corollary 3 this proves the claim.

Extending results of Mnëv [16] by using techniques described in [3] one finds (see [1, Cor. 9.5.11]) that there is a polynomial (Karp-)reduction of the problem to decide whether a system of linear inequalities has an integral solution to problem (S). Thus, problem (S) (and therefore, by Corollary 5, problem (G)) is not contained in $\text{co}\mathcal{NP}$, unless $\mathcal{NP} = \text{co}\mathcal{NP}$. Furthermore, there are rational simple polytopes P with the property that every rational simple polytope Q whose graph is isomorphic to the graph of P has vertices with super-polynomial coding lengths in the size of the graphs (this follows from Theorem B in [9]). Thus, it seems also unlikely that problem (G) is contained in $\mathcal{NP}$.

The results presented in Sect. 4 hence do neither lead to efficient algorithms nor to new examples of problems in $\mathcal{NP} \cap \text{co}\mathcal{NP}$ not (yet) known to be solvable in polynomial time. Nevertheless, they show that the problem to reconstruct a simple polytope from its graph can be modeled as a combinatorial optimization problem with a strongly dual problem. We hope that this is an appearance of Combinatorial Optimization Jack Edmonds is pleased to see in this volume dedicated to him.

7 Acknowledgements

I thank Günter M. Ziegler for valuable comments on an earlier version of the manuscript.

References

1. A. Björner, M. Las Vergnas, B. Sturmfels, N. White, and G. M. Ziegler. *Oriented Matroids (2nd ed.)*, volume 46 of *Encyclopedia of Mathematics and Its Applications*. Cambridge University Press, 1999.
2. R. Blind and P. Mani-Levitska. Puzzles and polytope isomorphisms. *Aequationes Math.*, 34:287–297, 1987.
3. J. Bokowski and B. Sturmfels. *Computational Synthetic Geometry*, volume 1355 of *Lecture Notes in Mathematics*. Springer, Heidelberg, 1989.
4. M. Develin. E-mail conversation, Nov 2000. develin@bantha.org.
5. J. Edmonds. Maximum matching and a polyhedron with 0,1-vertices. *J. Res. Natl. Bur. Stand. – B (Math. and Math. Phys.)*, 69B:125–130, 1965.
6. J. Edmonds. Paths, trees, and flowers. *Can. J. Math.*, 17:449–467, 1965.

7. M. R. Garey and D. S. Johnson. *Computers and Intractability. A Guide to the Theory of NP-Completeness*. W. H. Freeman and Company, New York, 1979.
8. M. X. Goemans and D. P. Williamson. The primal-dual method for approximation algorithms and its application to network design problems. In D. Hochbaum, editor, *Approximation Algorithms*, chapter 4. PWS Publishing Company, 1997.
9. J. E. Goodman, R. Pollack, and B. Sturmfels. The intrinsic spread of a configuration in $\mathbb{R}^d$. *J. Am. Math. Soc.*, 3(3):639–651, 1990.
10. C. Haase and G. M. Ziegler. Examples and counterexamples for Perles’ conjecture. Technical report, TU Berlin, 2001. To appear in: *Discrete Comput. Geometry*.
11. P. L. Hammer, B. Simeone, T. M. Liebling, and D. de Werra. From linear separability to unimodality: A hierarchy of pseudo-boolean functions. *SIAM J. Discrete Math.*, 1:174–184, 1988.
12. M. Joswig, V. Kaibel, and F. Körner. On the k -systems of a simple polytope. Technical report, TU Berlin, 2001. arXiv: math.CO/0012204, to appear in: *Israel J. Math.*
13. M. Joswig and G.M. Ziegler. Neighborly cubical polytopes. *Discrete Comput. Geometry*, 24:325–344, 2000.
14. V. Kaibel and M. Pfetsch. Computing the face lattice of a polytope from its vertex-facet incidences. Technical report, TU Berlin, 2001. arXiv:math.MG/01060043, submitted.
15. G. Kalai. A simple way to tell a simple polytope from its graph. *J. Comb. Theory, Ser. A*, 49(2):381–383, 1988.
16. N. E. Mnëv. The universality theorems on the classification problem of configuration varieties and convex polytopes varieties. In O.Ya. Viro, editor, *Topology and Geometry – Rohlin Seminar*, volume 1346 of *Lecture Notes in Mathematics*, pages 527–543. Springer, Heidelberg, 1988.
17. W. T. Tutte. A short proof of the factor theorem for finite graphs. *Can. J. Math.*, 6:347–352, 1954.
18. H. Whitney. Congruent graphs and the connectivity of graphs. *Am. J. Math.*, 54:150–168, 1932.
19. G. M. Ziegler. *Lectures on Polytopes*. Springer-Verlag, New York, 1995. Revised edition 1998.